

Version with markings to show changes made

In the claims:

A marked-up version of claims 132, 139, 140, and 141 is presented below.

132. (Twice Amended) A method for inducing myelination of a neural cell by a glial cell, said method comprising contacting said glial cell with an amount of a polypeptide which comprises an epidermal growth factor-like domain, wherein said epidermal growth factor-like domain comprises an amino acid sequence which is identical to an amino acid sequence encoded by a GGF/p185 erb B2 ligand gene, and wherein said amino acid sequence comprises an amino acid sequence encoded by a nucleic acid sequence selected from the group consisting of:

[SEQ ID NO: 154 (EGFL1);]

SEQ ID NO: 155 (EGFL2);

SEQ ID NO: 156 (EGFL3);

SEQ ID NO: 157 (EGFL4);

SEQ ID NO: 158 (EGFL5);

SEQ ID NO: 159 (EGFL6); and

amino acids 54-103 encoded by SEQ ID NO: 150.

139. (Twice Amended) The method of claim 132 [, 136, 137] or 141, wherein said method further comprises contacting said cell with a polypeptide which binds the p185 erb B2 receptor.

140. (Twice Amended) The method of claim 132 [, 136, 137] or 141, wherein said polypeptide is a recombinant polypeptide with glial cell mitogenic activity.

141. (Twice Amended) A method for inducing myelination of a neural cell by a glial cell, said method comprising contacting said glial cell with an amount of a polypeptide which comprises an epidermal growth factor-like domain, wherein said epidermal growth factor-like domain comprises an amino acid sequence which is identical to an amino acid sequence encoded by a GGF/p185 erb B2 ligand gene, and wherein said amino acid sequence comprises the amino acid sequence provided in [is selected from the group consisting of:

SEQ ID NO: 151;]

SEQ ID NO: 152 [; and

amino acids 362-411 of SEQ ID NO: 170],

wherein said contacting with said polypeptide is sufficient to induce myelination of said neural cell by said glial cell.

132. (Amended) A method for inducing myelination of a neural cell by a glial cell, said method comprising contacting said glial cell with an amount of a polypeptide which comprises an epidermal growth factor-like domain, wherein said epidermal growth factor-like domain comprises an [the] amino acid sequence [of] which is identical to an amino acid sequence encoded by a GGF/p185 erb B2 ligand gene, and wherein said amino acid sequence comprises an amino acid sequence encoded by a nucleic acid sequence selected from the group consisting of:

SEQ ID NO: 154 (EGFL1):

SEQ ID NO: 155 (EGFL2):

SEQ ID NO: 156 (EGFL3):

SEQ ID NO: 157 (EGFL4):

SEQ ID NO: 158 (EGFL5):

SEQ ID NO: 159 (EGFL6); and

amino acids 54-103 encoded by SEQ ID NO: 150

[sufficient to induce myelination of a neural cell by said glial cell].

136. (Amended) [The method of claim 133] A method for inducing myelination of a neural cell by a glial cell, said method comprising contacting said glial cell with an amount of a polypeptide which comprises an epidermal growth factor-like domain,

wherein said epidermal growth factor like domain [further] comprises the polypeptide encoded by SEQ ID NO: 188, wherein the human C/D'-segment polypeptide encoded by SEQ ID NO: 179 [178] [, wherein SEQ ID NO: 178] is immediately C-terminal to the human C-segment polypeptide encoded by SEQ ID NO: 177.

Sub D29 cont
C7

137. (Amended) [The method of claim 133] A method for inducing myelination of a neural cell by a glial cell, said method comprising contacting said glial cell with an amount of a polypeptide which comprises an epidermal growth factor-like domain, wherein said epidermal growth factor like domain [further] comprises the polypeptide encoded by SEQ ID NO: 189 [SEQ ID NO: 179], wherein the bovine C/D'-segment polypeptide encoded by SEQ ID NO: 143 [42] is immediately C-terminal to the human C-segment polypeptide encoded by SEQ ID NO: 177, and the human D-segment polypeptide encoded by SEQ ID NO: 180 is immediately C-terminal to the polypeptide encoded by SEQ ID NO: 143]

139. (Amended) The method of claim 132, 136, 137 or 141, wherein said [A] method [for inducing myelination of a neural cell by a glial cell,] further [comprising] comprises contacting said cell with [an amount of] a polypeptide which binds the p185 erb B2 receptor[, sufficient to induce myelination of a neural cell by said glial cell].

C⁷ 140. (Amended) The method of claim 132, 136, 137 or 141, wherein said polypeptide is [A method of inducing myelination of a neural cell by a glial cell, comprising contacting said glial cell with an amount of] a recombinant polypeptide with glial cell mitogenic activity [sufficient to induce myelination of a neural cell by said glial cell].

Add new claims 141-143.

C⁸ --141. A method for inducing myelination of a neural cell by a glial cell, said method comprising contacting said glial cell with an amount of a polypeptide which comprises an epidermal growth factor-like domain, wherein said epidermal growth factor-like domain comprises an amino acid sequence which is identical to an amino acid sequence encoded by a GGF/p185 erb B2 ligand gene, and wherein said amino acid sequence is selected from the group consisting of:
SEQ ID NO: 151;
SEQ ID NO: 152; and
amino acids 362-411 of SEQ ID NO: 170,
wherein said contacting with said polypeptide is sufficient to induce myelination of said neural cell by said glial cell.